

# **Fire Suppression & Native Range Release**

**Joseph G. Vollmer, Ph.D., BASF**



**Fire modeling conducted by:  
SYNERGY Resource Solutions Inc.**

**&**

**Special thanks to Kathy Kershaw  
of the BLM Lower Snake River  
District for her assistance.**

## Biomass data indicative of fine fuel loading from a BLM study area east of Boise

Type of Fuel	'96	'97	'98	'99	'00
	---- lbs/acre ----				
<i>Bromus</i>	1700	2222	2200	2245	1900
Litter	1500	2447	1350	3377	4182
Forbs	300	250	463	30	83
<b>Total</b>	<b>3500</b>	4919	4013	5652	<b>6255</b>

Source; 2001 Idaho Emergency Use Exemption, Plateau for control of leafy spurge, annual brome and medusahead in range. Idaho Department of Agriculture

# Fire Suppression & Native Range Release

- Study Plans & Results
- Fire Modeling – reducing fire risk.
- Native Range Release.
- Revegetation of *Bromus* infested areas.



# Methods used for the following research results

- 20 GPA
- 11002 flat fan tips
- CO<sup>2</sup> back pack sprayer
- 34 psi
- Small plot research  
10' x 50' w/ 3 reps




## Release of desirable plants

- BLM, Boise area, Point Addendum
- Area had moderate leaf litter layer w/ some desirable species present.
  - 90% *B. tectorum*
  - 10% WY big sage, Rabbit brush, Bottlebrush squirreltail, & Sandberg's bluegrass
- Treatments applied:
  - November 3, 1999



# Downy brome control

## Release of desirable plants



- **Application:**
  - Nov 3, 1999
  - Pre-emergence to downy brome germination
- **Rating:**
  - Spring = May, 2000
  - Fall = Nov, 2000
  - Spring = April 2001
  - Spring = April 2002





# Downy brome control

## Release of desirable plants



Non-treated check



May 3, 2000  
6 months after treatment  
Plateau 8oz/acre + MSO





# **Downy brome control**

## **Release of desirable plants**



**Area released from downy brome  
1½ years after treatment, Plateau 8oz/acre**

# Fire Modeling Data Collection




June 15, 2002 - 32 MAT

- plots were analyzed to determine fire behavior
- Data collected:
  - Biomass production
  - Litter accumulation
  - Plant height
  - Plant inter-space
- Data applied to:  
**BehavePlus** Fire-Modeling Program

# Fire Modeling – Live Biomass


Dry year – Tallest cheatgrass was 6 inches

Plant sample weights (grams/2 meter<sup>2</sup>)




# Fire Modeling – Biomass w/ Litter

Plant sample weights (grams/2 meter<sup>2</sup>)



# Fire Modeling

## Rate of Fire Spread




Rate of spread (m/min) at three slopes and three wind speeds and 5% fine fuel moisture comparing Plateau application rates of 0, 6, and 12 oz/ac.

Data collected and analyzed by Synergy Resource Solutions, Inc.



# Fire Modeling Flame Length



- flame length of 1.2 m (hand control ceases to be effective).
- - - flame length of 2.4 m (the fire cannot be contained from the head fire).


# Native Range Release Sheridan, WY

Non-Treated

Plateau 8 oz/A

Cheatgrass

Western wheatgrass






# **GRASS ESTABLISHMENT**

## **Boisie, ID BLM – Simco Road Site**

- **Treated: 11/3/99**
- **Planted: 11/17/99**
- **Timing: Pre-Plant, Pre-emergence to burbuttercup and downy brome.**
- **Species Planted:**
  - **Bozoisky Russian Wildrye**
  - **Vavilov Siberian Wheatgrass**




30 MAT Plateau applied after burn prior to planting of  
wildrye and wheatgrass

# DOWNY BROME - % CONTROL


## Boisie, ID – 30 MAT

5/21/02 (Pre-Plant)



# ESTABLISHMENT OF SIBERIAN WHEATGRASS & RUSSIAN WILD RYE - Boisie, ID – 30 MAT

# of Plants per 2 foot of row 5/21/02 (Pre-Plant)





# Revegetation after a burn



**Plateau 4oz  
1½ years after appl.**

**Plateau 2oz  
1½ years after appl.**

# Wyoming can still be rescued!










# Returning the Range



## Conclusion 3 YAT

- Treated areas had significantly less downy brome than non-treated area.
- Plateau at 6 oz/A or higher would effectively reduce *B. tectorum* in and around important habitat resources.
- Flame lengths would be reduced by 50 – 90% in the 6 oz rate of Plateau.

## **Conclusion 3 YAT**

- Plateau plots had Lower Flame Heights & Rate of Spread than the non-treated plots.
- Hand Control of fire could have been used on fires in all Plateau plots but not in any of the Non-treated plots.
- Plateau can be used to establish desired vegetation.

## **Conclusion 3 YAT**

- Plateau can be incorporated into land management plans, reducing the risk of loss of life, structures and vegetation in areas of concern by reducing fuel load.

## Conclusion

- Plateau is effective for control of annual *bromus spp.*
- Most rangeland grasses, forbs & shrubs are tolerant at these rates and timings.



# Annual Brome Control



**GAINING NEW GROUND.**